Sample



Emergency Care Plan

BEE STING ALLERGY

Student:	Grade	e: School Con	tact:		DOB:
	No (increased risk for se				
Father:		FHome #:	FWork #:		FCell #:
Emergency Contact:		Relationship	;	Phone:	
 MOUTH THROAT SKIN STOMACH LUNG HEART Th 	ALLERGIC REACTION Itching & swelling of lipe Itching, tightness in the Hives, itchy rash, swelling Nausea, abdominal crare Shortness of breath, reperturber of symptomic severity of symptomic important that treatments.	os, tongue or mouth oat, hoarseness, coughing of face and extremitings, vomiting, diarrhea petitive cough, wheezing out" ns can change quick	es S ly –	F THESE:	Student Photo
STAFF MEMBERS I	NSTRUCTED: ☐ Administration	☐ Classroom Teacher☐ Support Staff	` '	Special Area T Transportation	
TREATMENT: Remove stinger if visible, apply ice to area. Rinse contact area with water. Treatment should be initiated with symptoms without waiting for symptoms Benadryl ordered: Senadryl per provider's orders Call school nurse. Call parent/guardian if off school grounds. Epinephrine ordered: Senadryl per provider's orders If ANY SYMPTOMS BEYOND REDNESS OR SWELLING AT THE SITE OF THE STING ARE PRESENT AND EPINEPHRINE IS ORDERED, GIVE EPINEPHRINE IMMEDIATELY AND CALL 911. Preferred Hospital if transported: Epinephrine provides a 20 minute response window. After epinephrine, a student may feel dizzy or have an increased heart rate. This is a normal response. Students receiving epinephrine should be transported to the hospital by ambulance. A staff member should accompany the student to the emergency room if the parent, guardian or emergency contact is not present and adequate supervision for other students is present. Transportation Plan: Medication available on bus Medication NOT available on bus Does not ride bus Special instructions:					
Healthcare Provider:	23		Ph -		
Written by:	☐ Copy provided to Pare	nt 🔲 Cop	Dai v sent to Healt	e: hcare Provider	
Parent/Guardian Sign					